

## **Tumorigenic effects of dichloroacetic acid in female F344 rats**

Charles E. Wood<sup>1</sup>, Dennis E. House<sup>1</sup>, Michael H. George<sup>1</sup>, Stephen Kilburn<sup>1</sup>, Tanya M. Moore<sup>1</sup>, Anthony B. DeAngelo<sup>1</sup>

<sup>1</sup> National Health and Environmental Effects Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, NC

Corresponding author: Charles E. Wood, National Health and Environmental Effects Research Laboratory, U.S. Environmental Protection Agency, B105-03, 109 T.W. Alexander Drive, Research Triangle Park, NC 27711, USA, Email: wood.charles@epa.gov

**Introduction:** Dichloroacetic acid (DCA) is a halogenated organic acid produced during oxidant disinfection of drinking water. Prior studies indicate that DCA may increase liver tumors in mice. Here we evaluated the hepatic tumorigenicity of DCA in female rats when given alone or prior to phenobarbital (PB), a liver tumor promoter in rodents.

**Methods:** This study followed a stop-promotion design in which female F344 rats were divided into six treatment groups: (1) deionized water (dH<sub>2</sub>O, control); (2) dH<sub>2</sub>O switching to PB at 4 weeks; (3) DCA (1.0g/L); (4) DCA (1.5g/L) switching to dH<sub>2</sub>O at 4 weeks; (5) DCA (1.5g/L) switching to PB at 4 weeks; and (6) DCA (1.0g/L) switching to dH<sub>2</sub>O at 26 weeks (n=40-60/group). Treatment lasted for 100 weeks with interim pathology evaluations at week 52 (n=10/group for groups 1-5).

**Results:** No significant treatment-related effects on tumor incidence were observed for liver, adrenal gland, pituitary gland, uterus, or thyroid gland. Incidence and number of mammary gland fibroadenomas were higher in group 6 ( $p < 0.05$  vs. control). Incidence of preneoplastic hepatic lesions was greater for groups 2 and 5 (eosinophilic foci) and group 3 (clear cell foci) at 100 weeks ( $p < 0.05$  vs. control). Non-neoplastic treatment effects included chronic progressive nephropathy (CPN), hepatocyte single-cell necrosis, and centrilobular hepatocyte hypertrophy at 52 weeks (groups 2 and 5) and CPN at 100 weeks (groups 2-5) ( $p < 0.05$  vs. control).

**Discussion:** Chronic DCA exposure did not increase hepatic tumorigenesis in female rats. Further work is needed to confirm DCA effects on mammary gland tumors and CPN.

EPA Study No. 96-03  
PAI Work Assignment 96-11  
CSS Task 2.2.1, SSWR Task 2.2D

**Disclaimer:** The research described in this article has been reviewed by the National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency, and approved for publication. Approval does not signify that the contents necessarily reflect the views and the policies of the Agency, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.